CORRECTION Open Access



Correction: Susceptibility of AutoML mortality prediction algorithms to model drift caused by the COVID pandemic

Simone Maria Kagerbauer^{1,2*†}, Bernhard Ulm^{1,2†}, Armin Horst Podtschaske¹, Dimislav Ivanov Andonov¹, Manfred Blobner^{1,2}, Bettina Jungwirth² and Martin Graessner^{1,2}

Correction: Kagerbauer et al. BMC Medical Informatics and Decision Making (2024) 24:34

https://doi.org/10.1186/s12911-024-02428-z

Following the publication of the original article [1], the authors reported typesetting errors and a typo.

The first typesetting error was found in the Methods' section of the Abstract. The numbering in the section was mistakenly linked to the reference, as follows:

We trained different ML models with the H2O AutoML method on a dataset comprising 102,666 cases of surgical patients collected in the years 2014–2019 to predict post-operative mortality using preoperatively available data. Models applied were Generalized Linear Model with regularization, Default Random Forest, Gradient Boosting Machine, eXtreme Gradient Boosting, Deep Learning and Stacked Ensembles comprising all base models. Further, we modified the original models by applying three different methods when training on the original

pre-pandemic dataset: (Rahmani K, et al., Int J Med Inform 173:104930, 2023) we weighted older data weaker, (Morger A, et al., Sci Rep 12:7244, 2022) used only the most recent data for model training and (Dilmegani C, 2023) performed a z-transformation of the numerical input parameters. Afterwards, we tested model performance on a pre-pandemic and an in-pandemic data set not used in the training process, and analysed common features.

The correct sentence should have been:

We trained different ML models with the H2O AutoML method on a dataset comprising 102,666 cases of surgical patients collected in the years 2014–2019 to predict postoperative mortality using preoperatively available data. Models applied were Generalized Linear Model with regularization, Default Random Forest, Gradient Boosting Machine, eXtreme Gradient Boosting, Deep Learning and Stacked Ensembles comprising all base models. Further, we modified the original models by applying three different methods when training on the original pre-pandemic dataset: (1) we weighted older data weaker, (2) used only the most recent data for model training and (3) performed a z-transformation of the numerical input parameters. Afterwards, we tested model performance on a pre-pandemic and an in-pandemic data set not used in the training process, and analysed common features.

The second error was found in the legend of Fig. 4, following the abbreviation of ASA which mentions the

The online version of the original article can be found at https://doi.org/10.1186/s12911-024-02428-z.

*Correspondence: Simone Maria Kagerbauer simone.kagerbauer@uni-ulm.de

Germany

¹Department of Anaesthesiology and Intensive Care Medicine, School of Medicine, Technical University of Munich, Munich, Germany ²Department of Anaesthesiology and Intensive Care Medicine, School of Medicine, University of Ulm, Albert-Einstein-Allee 23, 89081 Ulm,



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

[†]Simone Maria Kagerbauer and Bernhard Ulm are shared first authors of the manuscript.

number 34. It reads: ASA: American Society of Anaesthesiologists Physical Score 34.

The correct explanation should have referred to the source of Mayhew D, Mendonca V, Murthy BVS. A review of ASA physical status- historical perspectives and modern developments. Anaesthesia 2019;74:373–9. https://doi.org/10.1111/anae. 14,569. Thus, it should have read: ASA: American Society of Anaesthesiologists Physical Score [16].

Moreover, there is a typo with regards to the first line of the legend of Table 2 where the word "pandemic" was repeated twice. The correct legend should have read "The Kolmogorov-Smirnov-Test was performed between the pre-pandemic validation and the pre-pandemic test set". The original article [1] has been updated.

Published online: 19 February 2024

References

 Kagerbauer SM, Ulm B, Podtschaske AH, et al. Susceptibility of AutoML mortality prediction algorithms to model drift caused by the COVID pandemic. BMC Med Inf Decis Mak. 2024;24:34. https://doi.org/10.1186/ s12911-024-02428-z.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.