

Research article

Clinical outcomes resulting from telemedicine interventions: a systematic review

William R Hersh^{*1,4}, Mark Helfand^{1,4}, James Wallace^{1,4}, Dale Kraemer^{1,4}, Patricia Patterson^{1,2,4}, Susan Shapiro^{2,4} and Merwyn Greenlick^{3,4}

Address: ¹Division of Medical Informatics & Outcomes Research, Oregon Health & Science University, BICC, 3181 SW Sam Jackson Park Rd, Portland, OR 97201, USA, ²School of Nursing, Oregon Health & Science University, BICC, 3181 SW Sam Jackson Park Rd, Portland, OR 97201, USA, ³Department of Public Health & Preventive Medicine, Oregon Health & Science University, BICC, 3181 SW Sam Jackson Park Rd, Portland, OR 97201, USA and ⁴Evidence-Based Practice Center, Oregon Health & Science University, BICC, 3181 SW Sam Jackson Park Rd, Portland, OR 97201, USA

E-mail: William R Hersh* - hersh@ohsu.edu; Mark Helfand - helfand@ohsu.edu; James Wallace - wallacej@ohsu.edu; Dale Kraemer - kraemerd@ohsu.edu; Patricia Patterson - patterso@ohsu.edu; Susan Shapiro - shapiros@ohsu.edu; Merwyn Greenlick - mitchg@ohsu.edu

*Corresponding author

Published: 26 November 2001

Received: 23 October 2001

BMC Medical Informatics and Decision Making 2001, 1:5

Accepted: 26 November 2001

This article is available from: <http://www.biomedcentral.com/1472-6947/1/5>

© 2001 Hersh et al; licensee BioMed Central Ltd. Verbatim copying and redistribution of this article are permitted in any medium for any non-commercial purpose, provided this notice is preserved along with the article's original URL. For commercial use, contact info@biomedcentral.com

Abstract

Background: The use of telemedicine is growing, but its efficacy for achieving comparable or improved clinical outcomes has not been established in many medical specialties. The objective of this systematic review was to evaluate the efficacy of telemedicine interventions for health outcomes in two classes of application: home-based and office/hospital-based.

Methods: Data sources for the study included reports of studies from the MEDLINE, EMBASE, CINAHL, and HealthSTAR databases; searching of bibliographies of review and other articles; and consultation of printed resources as well as investigators in the field. We included studies that were relevant to at least one of the two classes of telemedicine and addressed the assessment of efficacy for clinical outcomes with data of reported results. We excluded studies where the service did not historically require face-to-face encounters (e.g., radiology or pathology diagnosis). All included articles were abstracted and graded for quality and direction of the evidence.

Results: A total of 25 articles met inclusion criteria and were assessed. The strongest evidence for the efficacy of telemedicine in clinical outcomes comes from home-based telemedicine in the areas of chronic disease management, hypertension, and AIDS. The value of home glucose monitoring in diabetes mellitus is conflicting. There is also reasonable evidence that telemedicine is comparable to face-to-face care in emergency medicine and is beneficial in surgical and neonatal intensive care units as well as patient transfer in neurosurgery.

Conclusions: Despite the widespread use of telemedicine in virtually all major areas of health care, evidence concerning the benefits of its use exists in only a small number of them. Further randomized controlled trials must be done to determine where its use is most effective.

Background

There are over 450 telemedicine programs worldwide, with over 360 of those in the United States [1]. There are

programs in virtually every medical specialty, and the populations they most commonly serve include those who live in rural areas, the elderly, and veterans. With

the growing ability of modern computer and communications technology to capture and quickly transmit textual, audio, and video information, many have advocated its use to improve the health care provided to individuals in rural areas, in the home, and in other places where medical personnel are not readily available.

Many believe that the growth of telemedicine is impeded by reimbursement policies of the federal government and private insurers [2,3]. These authors claim this leads to a vicious cycle: There is a growing call for telemedicine services to be covered as part of health insurance. Health care payors, whether governmental or private, have been reluctant to do so. This, in turn, has led telemedicine advocates to blame payors for impeding development of the technology.

One reason for the lack of coverage of telemedicine has been an uncertainty about its efficacy and cost. There have been a number of previous systematic reviews assessing the efficacy of telemedicine and related technologies [4–9]. All of these studies noted that although the technology showed promise in certain areas, the overall methodologic quality of the evaluative studies was low and the plan for the most appropriate and cost-effective use of telemedicine was unclear.

This systematic review had its origins in a report commissioned by the US Health Care Financing Agency (HCFA) and the Agency for Healthcare Research and Quality (AHRQ) to assess the efficacy of telemedicine in the adult, non-pregnant population [1]. A subsequent supplemental report extended the analysis to the pediatric and obstetric population [10]. The goal of this paper is to assess whether telemedicine interventions result in comparable health outcomes to in-person care. It merges results from the original two reports and updating them with studies published since the reports were submitted to the federal agencies.

We broadly divide telemedicine into two general areas. In *office/hospital-based telemedicine*, both the patient and clinician are in a professional health care setting, such as a practitioner's office or in the hospital. There is a subcategory of office/hospital-based telemedicine called *store-and-forward telemedicine*, where clinical data is collected and forwarded for review later in a completely asynchronous manner, but it was not assessed in this systematic review because our literature review failed to identify any studies of clinical outcomes utilizing it. In *home-based telemedicine*, the patient is at his or her residence, with direct communication between the clinician and the patient or their caregiver.

Initial search strategy

- 1 exp telemedicine/
- 2 telemedicine.mp.
- 3 telehealth.tw.
- 4 remote consultation\$.mp.
- 5 1 or 2 or 3 or 4
- 6 exp home care services/
- 7 home nursing/
- 8 6 or 7
- 9 exp therapy, computer-assisted/
- 10 exp computers/
- 11 exp computer communication networks/
- 12 exp medical informatics/
- 13 exp telecommunications/
- 14 exp monitoring, physiologic/
- 15 monitor\$.tw.
- 16 blood glucose self-monitoring/
- 17 self-examination/
- 18 self exam\$.tw.
- 19 self monitor\$.tw.
- 20 self test\$.tw.
- 21 14 or 15 or 16 or 17 or 18 or 19 or 20
- 22 tele\$.tw.
- 23 (remote or offsite or distance).tw. tw=abstract, title
- 24 rural population/
- 25 rural health services/
- 26 hospitals, rural/
- 27 rural.tw.
- 28 22 or 23 or 24 or 25 or 26 or 27
- 29 21 and 28
- 30 9 or 10 or 11 or 12 or 13 or 29
- 31 8 and 30
- 32 31 not 5
- 33 limit 32 to english language
- 34 32 not 33
- 35 limit 34 to abstracts
- 36 33 or 35

Supplemental Search Strategy (to identify more home-based telemedicine articles)

- 1 exp computer communication networks
- 2 patient participation
- 3 exp consumer satisfaction
- 4 delivery of health care
- 5 exp home care services
- 6 exp home nursing
- 7 house calls/ or housecalls.mp
- 8 2 or 3 or 4 or 5 or 6 or 7
- 9 1 and 8
- 10 limit 9 to english language

Figure 1

Search strategies. Search strategies for MEDLINE shown; comparable strategies were used for EMBASE, CINAHL, and HealthSTAR.

Methods

The criteria for study inclusion in this systematic review were that the study be relevant to at least one of the areas of telemedicine and contain data on outcomes of clinical care that compared telemedicine with a control group of in-person care. We excluded services that would not normally require face-to-face encounters between the clinician and patient (e.g., radiology and pathology diagnosis), used only telephone care or electronic mail, or provided medical advice directly to the public.

Table 1: Classification of evidence for studies of clinical outcomes of telemedicine interventions.

| Study Class | Characteristic |
|-------------|---|
| I | <ul style="list-style-type: none"> • Properly designed random controlled trials |
| II | <ul style="list-style-type: none"> • Random controlled trials that contain design flaws preventing specification of Class I • Properly designed trials with control groups not randomized • Multi-center or population-based longitudinal (cohort) study • Case control studies |
| III | <ul style="list-style-type: none"> • Descriptive studies (uncontrolled case series) • Clinical experience • Expert opinion • Case reports |

We developed a search strategy designed to find any publications about telemedicine and used it to search MEDLINE, EMBASE, CINAHL, and HealthSTAR through February, 2001 (see Figure 1). We also searched through telemedicine reports and compilations such as the *Conseil d'Evaluation des Technologies de la Sante du Quebec*[11], the *Telemedicine Strategic Healthcare Group Report*,[12] the *International Society for Telemedicine Conference Proceedings*, the Association of Telemedicine Service Providers' database [13], and the *Telemedicine Sourcebook 1998*[14]. We also assessed three systematic reviews (different in scope from this study) from the International Network of Agencies for Health Technology (INHATA) [5], the Cochrane Database of Systematic Reviews [7], and the Agenda d'Aval-uacio de Tecnologia Medica [6]. In addition, we identified additional articles from the reference lists of included reports and articles and also contacted known telemedicine experts to find additional articles to identify and describe telemedicine programs. Finally, we also handsearched all issues of the two major telemedicine journals, *Journal of Telemedicine and Telecare* and *Telemedicine Journal* (recently renamed *Telemedicine Journal and e-Health*), through the end of 2000.

Each title/abstract retrieved was reviewed by two reviewers. When the two reviewers disagreed, a third reviewer made the final decision. We retrieved the full-text articles for citations selected for possible inclusion in the systematic review. When an article met the criteria for

inclusion, the summary statistics were extracted. To evaluate the quality of the studies assessed, we adapted a scale developed by our evidence-based practice center (Table 1) [15]. We used a second scale to indicate the direction of the evidence (Table 2).

Results

Literature searching in MEDLINE, CINAHL, EMBASE, and HealthSTAR yielded 4,628 possibly relevant references. Additional studies were identified from previous systematic reviews, reference lists of included papers, and handsearching of the two peer-reviewed telemedicine publications. Applying the inclusion criteria yielded 19 articles in home-based telemedicine (Table 3) and six in interactive office/hospital-based telemedicine (Table 4).

The overall methodologic quality of many of the studies was low. Only eight home-based telemedicine studies and one office/hospital-based telemedicine studies met the criteria for class I studies. Some of the problems included small sample sizes, short duration periods for the interventions, and lack of follow-up after the initial episode of care. For this reason, as well as the general heterogeneous nature of the studies, no attempt at aggregation (i.e., meta-analysis) was performed.

Home-based telemedicine

By far the most common area that has been studied in home-based telemedicine is monitoring of blood sugar in patients with diabetes mellitus, with a total of eight studies. The next most common areas have included general chronic disease management (three studies), hypertension (two studies), and AIDS (two studies).

Diabetes Mellitus

The studies that have assessed home monitoring of diabetes have focused on tracking insulin dosages and used Hemoglobin A 1C as an outcome measure. Two of the studies identified focused on gestational diabetes, one on children, and the remainder on adults. Only one diabetes study was adequately designed to rate as a class I randomized controlled trial (RCT) [16]. This pediatric study showed no difference in glycemic control between

Table 2: Direction of effect for evidence

| Study Class | Characteristic |
|-------------|---|
| A | Strong positive effect |
| B | Weak positive effect |
| C | Conflicting evidence for effect |
| D | Negative effect (evidence that the technology is inferior or ineffective) |

Table 3: Studies of clinical outcomes using interventions of home-based telemedicine.

| Outcomes | Quality Score | Clinical Specialty | Sample | Intervention | Effects |
|---------------------|---------------|---------------------|----------------|---|--|
| Flatley-Brennan[29] | I-B | AIDS | 57 patients | Social isolation and decision-making skill for home computer network (RCT) | Reduced social isolation when controlled for depression and improved confidence in decision-making with increased use |
| Gustafson[30] | I-A | AIDS | 183 patients | Quality of life and hospitalizations (RCT) | Improvement in active life, negative emotions, cognitive functions, social support, and participation in health care; fewer hospitalizations |
| Brennan[31] | I-B | Alzheimer's Disease | 102 caregivers | Social isolation and decision-making skill for home computer network (RCT) | Improved decision-making confidence but no improvement in decision-making skill or social isolation |
| Sparks [32] | II-B | Cardiology | 20 patients | Comparison of home exercise program with transtelephonic exercise monitoring vs. hospital-based program (RCT) | Both groups improved equally in cardiac function, no medical emergencies in either group |
| Mahmud [26] | III-B | Chronic Disease | 12 patients | Home telecare in chronic disease for frail elderly | Improved compliance and control of disease process; decreased hospitalization and nursing home placement |
| Nakamura[25] | II-B | Chronic Disease | 32 patients | Home telecare in chronic disease for frail elderly | Improvement in activities of daily living, communication, and social cognition |
| Johnston[24] | I-B | Chronic Disease | 212 patients | Home telecare in chronic disease for frail elderly (RCT) | Both groups had comparable medication compliance, knowledge of disease, and ability for self-care |
| Ahring[17] | II-B | Diabetes Mellitus | 42 patients | Home blood sugar monitoring (RCT) | Computer group had HgbA1c drop from 10.6% to 9.2% (-13.2%); control group from 11.2% to 10.2% (-8.9%) |
| Shultz[21] | II-B | Diabetes Mellitus | 20 patients | Home blood sugar monitoring (RCT) | Reduced HgbA1c levels in computer group but details not given |
| Billiard[19] | II-B | Diabetes Mellitus | 22 patients | Home blood sugar monitoring (RCT) | Computer group had HgbA1c drop from 6.7% to 6.0%; control group from 6.8% to 6.7% |
| DiBiase[22] | II-B | Diabetes Mellitus | 20 patients | Home blood sugar monitoring in gestational diabetes (RCT) | Computer group had HgbA1c drop from 6.4% to 5.0%; control group from 7.1% to 5.7% |
| Frost[23] | II-B | Diabetes Mellitus | 21 patients | Home blood sugar monitoring in gestational diabetes | Computer group had HgbA1c drop from 6.1% to 5.4%; control group from 6.2% to 5.7% |
| Marrero[16] | I-B | Diabetes Mellitus | 106 patients | Home blood sugar monitoring (RCT) | Computer group had HgbA1c rise from 9.4% to 10.0%; control group from 9.9% to 10.3%; no difference in ER visits, psychological status, or family functioning |
| Mease[20] | II-B | Diabetes Mellitus | 28 patients | Home blood sugar monitoring (RCT) | Computer group had HgbA1c fall from 9.5% to 8.2% vs. 9.5% to 8.6% for control group |
| Biermann[18] | II-B | Diabetes Mellitus | 46 patients | Home blood sugar monitoring (RCT) | Computer group had HgbA1c fall from 8.3% to 7.3% vs. 8.0% to 6.8% for control group |
| Friedman[27] | I-A | Hypertension | 267 patients | Automated patient monitoring and counseling (RCT) | Adherence and diastolic blood pressure improved |
| Cartwright[28] | I-B | Hypertension | 99 patients | Anxiety, blood pressure readings, and gestational age at delivery in home vs. hospital-monitored women (RCT) | Comparable levels of anxiety, mean blood pressure, and gestational age of delivery |
| Gray[33] | I-B | Neonatology | 56 patients | Quality of care and hospitalization (RCT) | Trend towards earlier discharge from hospital |
| Miyasaka[34] | III-B | Pulmonary | 10 patients | Amount of unscheduled care before and after installation of videophone access to physician | Reduction in number of house calls (5 vs. 0), unscheduled hospital visits (24 vs. 5), and hospital admission days (22 vs. 10) |

Table 4: Studies of clinical outcomes using interventions of office/hospital-based telemedicine.

| Outcomes | Quality Score | Clinical Specialty | Sample | Intervention | Effects |
|---------------|---------------|--------------------|--------------|---|---|
| Wootton[37] | II-B | Dermatology | 204 patients | Need for special follow-up (RCT) | No difference in need for follow-up |
| Brennan[36] | I-A | Emergency Medicine | 100 patients | Patients randomized to local or telemedicine care (RCT) | No difference in ER return or need for additional care |
| Rosenfeld[38] | II-B | Intensive Care | 201 patients | Addition of remote intensivist to surgical ICU | Decreases in severity-adjusted ICU mortality (46–68%) and hospital mortality (30–33%). Decreases in ICU complications (44–50%) and ICU length of stay (30–34%). |
| Rendina[39] | II-B | Neonatology | 314 patients | Length of stay in NICU for telemedicine vs. no telemedicine | Length of stay decreased significantly related to birth weight |
| Goh[40] | III-B | Neurosurgery | 116 patients | Neurosurgery transfer before and after teleradiology | Fewer adverse events during transfer (8% vs. 32%) |
| Goh[41] | III-B | Neurosurgery | 63 patients | Head injury patients with teleradiology | Fewer adverse events during transfer (6.4% vs. 32.1%) |

groups (HgbA1C for both groups actually rose), emergency room visits, psychological status, and family functioning.

The two largest studies of adults showed that both the experimental and control groups improved, with no statistically significant difference between them [17,18]. Three other studies demonstrated a small but statistically significant benefit in HgbA1C [19–21].

Both studies of home gestational diabetes monitoring found improvements in blood sugar values but not in HgbA1C, although each had very small sample sizes and probable inadequate statistical power to detect a difference [22,23].

General Chronic Disease Management

Three studies addressed general chronic disease management. The largest and best-designed was an RCT where both the intervention and control groups were provided usual home health care but the intervention group also received a video system that allowed real-time remote interaction with the health system [24]. Both groups had comparable compliance with medication regimen, knowledge about their disease, ability to move toward self-care, and scores on the 12-item Short-Form Health Survey (SF-12)TM.

The other studies described the use of videophones for patient communication with health care providers. In a study in Japan, 16 elderly patients who were provided with videophones were compared to 16 matched patients who used regular home health services [25]. After 3 months, the videophone group significantly exceeded the control group in activities of daily living, communica-

tions, and social cognition, as measured with the Functional Independence Measure. The videophone intervention was a supplement to (not a substitute for) regular home health services. The second study, a qualitative report of 12 cases, showed a reduction in the number of home-care visits needed for seven of the patients [26].

Hypertension

Of the two studies of monitoring hypertension via home-based telemedicine, one looked at the elderly while the other looked at pregnant women. The study of the elderly was an RCT that assessed the effect of a computer-controlled, automated telephone system versus usual office-based care on adherence and blood pressure control in older hypertensive patients [27]. Mean antihypertensive medication adherence improved 17.7 percent for telephone system users versus 11.7 percent for controls. Mean diastolic blood pressure decreased 5.2 mm Hg in users compared to a 0.8 mm Hg drop in controls, a statistically significant difference. There was also a positive relationship between medication adherence and blood pressure reduction. The study of hypertension in pregnant women found that an at-home blood pressure monitoring system resulted in comparable levels of blood pressure, anxiety, and gestational age of delivery as hospital-based monitoring [28].

AIDS

Two studies of persons with AIDS demonstrated the value of a home computer link to information, others with the disease, and a health care professional who could steer the patient to advice. One of the studies showed the system reduced social isolation after controlling for depression as well as improved decision-making confi-

dence [29]. The other demonstrated various quality-of-life improvements as well as fewer and shorter hospitalizations [30].

Other areas

The remaining studies provided evidence of benefit in a variety of areas:

- An RCT of a home computer link for caregivers of patients with Alzheimer's Disease found improved decision-making confidence [31].
- An RCT of patients in a home-exercise program using transtelephonic exercise monitoring compared to a hospital-based program improved cardiac function by a similar amount [32]. (One caveat of this study was that it had such low statistical power that it would not be likely to detect any difference between the two interventions in efficacy or in complications even if they existed.)
- An RCT of an Internet-based telemedicine program for families of children in a neonatal intensive care unit found that among infants born with weight of <1,000 grams, there was a trend towards shorter hospital stays [33].
- A study of videophones for pediatric home ventilator patients found that the number of unscheduled hospital visits and hospital admission days was reduced significantly compared with historical controls [34].

Office/hospital-based telemedicine

The six studies assessing clinical outcomes from interventions of office/hospital-based telemedicine were spread across five specialties. (We actually identified another study that appeared to be a preliminary report of another study in the group [35].)

Of the two RCTs identified, one assessed outcomes in patients entering an emergency department who were randomized to in-person or telemedicine care [36]. There were no differences in the need for additional follow-up care or return to the emergency room (ER), showing that telemedicine was as effective as regular care in this setting. Another RCT assessed "clinical outcome," which was defined as the need to have a follow-up appointment with a hospital-based specialist [37]. Interactive teledermatology consultation (46 percent) was found to have the same rate of need for follow-up care as in-person consultation (45 percent). No statistical analysis of the differences was performed.

A time series cohort study demonstrated reduced mortality, complications, and length of stay in a community hospital surgical intensive care unit (ICU) with continu-

ous intensivist oversight via telemedicine [38]. A study in a neonatal ICU used a retrospective comparison to show reduced length of stay with the use of a telecardiology system [39]. Two studies of neurosurgery patients demonstrated benefit of transmission of radio logic studies to avert complications in patient transfer [40,41].

Discussion

Despite the widespread use of telemedicine in virtually all areas of health care delivery, there is only a small amount of evidence that interventions provided by telemedicine result in clinical outcomes are comparable to or better than face-to-face care. The best evidence comes from home-based telemedicine, where modest benefits have been shown for patients with chronic disease, AIDS, and Alzheimer's Disease. The most studied area in home-based telemedicine is monitoring of blood sugar in patients with diabetes mellitus. The benefits are not conclusive, with studies showing no change or a slight drop in HgbA1C levels.

In office/hospital-based telemedicine, there is good evidence that telemedicine interventions provide comparable care in the ER settings and benefit in the surgical and neonatal ICU settings. There is also probable benefit in its use prior to transfer of neurosurgery patients. Although there is good evidence for the diagnostic capability of teledermatology [1], the only assessment of its outcomes has focused on the need for specialty follow-up.

Further studies must be done to demonstrate the efficacy of telemedicine for diagnostic and management decisions. Large-scale RCTs must be done to identify the health outcomes whose benefit appears most promising. If the goal is to show comparability to usual care, then studies must provide adequate statistical power to show that the lack of a difference truly exists. Small studies with inadequate power are not good evidence. The fact that telemedicine is an emerging technology is not a reason for failing to perform appropriate evaluation studies. Rather, new methodologies such as "tracker trials" should be used to assess it systematically [42]. Tracker trials are designed to assess new and/or rapidly changing interventions and compare efficacy not only of the general intervention but also specific instances of it, such as a newly-developed approach that has become available after the general trial started. We would also advocate that journal editors exhibit restraint when considering publication of low-quality evaluation studies. As such studies often accompany otherwise appropriate descriptions of telemedicine systems, they should consider refusing to publish the portion of the article containing the low-quality evaluation study and publish the rest.

The growth of high-speed telecommunications networks and their ability to transmit higher-quality imaging and other clinical information indicates a bright future for telemedicine, especially as the infrastructure is built out to reach to individuals who have historically been denied high-quality health care due to their remote location. As the cost of these services will challenge the health care budgets of even well-to-do nations, it is imperative that decisions to use them be based on appropriate evidence that they lead to comparable patient outcomes in office/hospital-based telemedicine and improved patient outcomes in home-based telemedicine.

Disclaimer

The authors of this article are responsible for its contents, including any clinical or treatment recommendations. No statement in this article should be construed as an official position of the Agency for Healthcare Research and Quality or the U.S. Department of Health and Human Services.

Competing Interests

None declared.

Acknowledgments

This study was conducted by the OHSU Evidence-based Practice Center under Agency for Healthcare Research and Quality contract 290-97-0018, task order no. 3, Rockville, MD.

References

- Hersh WR, Helfand M, Wallace JA, Kraemer DK, Patterson PK, Shapiro SE, Greenlick MR, Chan BKS, Eilers GM: **Telemedicine for the Medicare Population. Evidence Report/Technology Assessment No. 24. AHRQ Publication No. 01-E012.** Rockville, MD: Agency for Healthcare Research and Quality; 2001
- Grigsby J, Sanders J: **Telemedicine: where it is and where it's going.** *Annals of Internal Medicine* 1998, **129**:123-127
- Strode SW, Gustke S, Allen A: **Technical and clinical progress in telemedicine.** *Journal of the American Medical Association* 1999, **281**:1066-1068
- Balas EA, Jaffrey F, Kuperman GJ, Boren SA, Brown GD, Pincioli F, Mitchell JA: **Electronic communication with patients evaluation of distance medicine technology.** *Journal of the American Medical Association* 1997, **278**:152-159
- Ohinmaa A, Hailey D, Roine D: **The Assessment of Telemedicine: General Principles and a Systematic Review.** Helsinki, Finland: Finnish Office for Health Care Technology Assessment; 1999
- Almazan C, Gallo P: **Assessing Clinical Benefit and Economic Evaluation in Telemedicine.** Barcelona, Spain: Catalan Agency for Health Technology Assessment; 1999
- Currell R, Urquhart C, Wainwright P, Lewis R: **Telemedicine versus face to face patient care: effects on professional practice and health care outcomes.** *Cochrane Database of Systematic Reviews [computer file]* 2000CD002098
- Mair F, Whitten P: **Systematic review of studies of patient satisfaction with telemedicine.** *British Medical Journal* 2000, **320**:1517-1520
- Whitten P, Kingsley C, Grigsby J: **Results of a meta-analysis of cost-benefit research: is this a question worth asking?** *Journal of Telemedicine & Telecare* 2000, **6**:S4-S6
- Hersh WR, Wallace JA, Kraemer DK, Patterson PK, Nichols P, Greenlick MR, Helfand M: **Telemedicine for the Medicare Population: Pediatric, Obstetric, and Clinician-Indirect Home Interventions. Evidence Report/Technology Assessment No. 24, Supplement. AHRQ Publication No. 01-E060.** Rockville, MD: Agency for Healthcare Research and Quality; 2001
- Anonymous: **Telehealth and Telemedicine in Quebec: Current Issues.** Quebec: Conseil d'Evaluation des Technologies de la Sante du Quebec; 1999
- Anonymous: **Issues and Opportunities Facing VHA Telemedicine: Reinventing Service To Patients.** Washington, DC: Department of Veteran's Affairs Telemedicine Strategic Healthcare Group; 1999
- Grigsby B, Brown N: **Report on U.S. Telemedicine Activity.** Portland, OR: Association of Telemedicine Service Providers; 1998
- Rosenthal BE: **Telemedicine Sourcebook 1998: A Resource Guide for the Practical Application of Telemedicine in the Managed Care Marketplace.** New York: Faulkner & Gray, Inc.; 1997
- Chesnut RM, Carney N, Maynard H, Patterson PK, Mann NC, Helfand M: **Rehabilitation for Traumatic Brain Injury. Evidence Report/Technology Assessment No. 2. AHRQ Publication No. 99-E006.** Rockville, MD: Agency for Health Care Policy and Research; 1999
- Marrero DG, Vandagriff JL, Kronz K, Fineberg NS, Golden MP, Gray D, Orr DP, Wright JC, Johnson NB: **Using telecommunication technology to manage children with diabetes: the Computer-Linked Outpatient Clinic (CLOC) Study.** *Diabetes Educator* 1995, **21**:313-319
- Ahring KK, Ahring JP, Joyce C, Farid NR: **Telephone modem access improves diabetes control in those with insulin-requiring diabetes.** *Diabetes Care* 1992, **15**:971-975
- Biermann E, Dietrich W, Standl E: **Telecare of diabetic patients with intensified insulin therapy: a randomized clinical trial.** *Studies in Health Technology & Informatics* 2000, **77**:327-332
- Billiard A, Rohmer V, Roques M, Joseph M, Suraniti S, Giraud P, Limal J, Fressinaud P, Marre M: **Telematic transmission of computerized blood glucose profiles for IDDM patients.** *Diabetes Care* 1991, **14**:130-134
- Mease A, Whitlock WL, Brown A, Moore K, Pavlisacsak H, Dingbaum A, Laceyfield D, Buker K, Xenakis S: **Telemedicine improved diabetic management.** *Military Medicine* 2000, **165**:579-584
- Shultz EK, Bauman A, Hayward M, Holzman R: **Improved care of patients with diabetes through telecommunications.** *Annals of the New York Academy of Sciences* 1992, **670**:141-145
- di Biase N, Napoli A, Sabbatini A, Borrello E, Buongiorno AM, Falluca F: **Telemedicine in the treatment of diabetic pregnancy.** *Annali dell Istituto Superiore di Sanita* 1997, **33**:347-351
- Frost D, Beischer W: **Telemedicine in the management of pregnancy in type I diabetic women [letter].** *Diabetes Care* 2000, **23**:863-864
- Johnston B, Wheeler L, Deuser J, Sousa KH: **Outcomes of the Kaiser Permanente Tele-Home Health Research Project.** *Archives of Family Medicine* 2000, **9**:40-45
- Nakamura K, Takano T, Akao C: **The effectiveness of videophones in home healthcare for the elderly.** *Medical Care* 1999, **37**:117-125
- Mahmud K, Lenz J: **The personal telemedicine system: a new tool for the delivery of health care.** *Journal of Telemedicine & Telecare* 1995, **1**:173-177
- Friedman RH, Kazis LE, Jette A, Smith MB, Stollerman J, Torgerson J, Carey K: **A telecommunications system for monitoring and counseling patients with hypertension. Impact on medication adherence and blood pressure control.** *American Journal of Hypertension* 1996, **9**:285-292
- Cartwright W: **Objective measurement of anxiety in hypertensive pregnant women managed in hospital and in the community.** *British Journal of Obstetrics and Gynaecology* 1992, **99**:182-185
- Flatley-Brennan P: **Computer network home care demonstration: a randomized trial in persons living with AIDS.** *Computers in Biology & Medicine* 1998, **28**:489-508
- Gustafson DH, Hawkins R, Boberg E, Pingree S, Serlin RE, Graziano F, Chan CL: **Impact of a patient-centered, computer-based health information/support system.** *American Journal of Preventive Medicine* 1999, **16**:1-9
- Brennan PF, Moore SM, Smyth KA: **The effects of a special computer network on caregivers of persons with Alzheimer's Disease.** *Nursing Research* 1995, **44**:166-172
- Sparks KE, Shaw DK, Eddy D, Hanigovsky P, Vantrese J: **Alternatives for cardiac rehabilitation patients unable to return to a hospital-based program.** *Heart & Lung* 1993, **22**:298-303
- Gray J, Safran C, Davis RB, Pompilio-Weitzner G, Stewart JE, Zaccagnini L, Pursley D: **Baby CareLink: using the Internet and**

- telemedicine to improve care of high-risk infants.** *Pediatrics* 2000, **106**:1318-1324
34. Miyasaka K, Suzuki Y, Sakai H, Kondo Y: **Interactive communication in high-technology home care: videophones for pediatric ventilatory care.** *Pediatrics* 1997, **99**:E1
 35. Loane M, Bloomer S, Corbett R, Eedy D, Hicks N, Lotery H, Mathews C, Paisley J, Steele K, Wootton R: **A randomized controlled trial to assess the clinical effectiveness of both realtime and store-and-forward teler dermatology compared with conventional care.** *Journal of Telemedicine and Telecare* 2000, **6**:S1-S3
 36. Brennan JA, Kealy JA, Gerardi LH, Shih R, Allegra J, Sannipoli L, Lutz D: **randomized controlled trial of telemedicine in an emergency department.** *Journal of Telemedicine & Telecare* 1998, **4**:18-20
 37. Wootton R, Bloomer SE, Corbett R, Eedy DJ, Hicks N, Lotery HE, Mathews C, Paisley J, Steele K, Loane MA: **Multicentre randomised control trial comparing real time teler dermatology with conventional outpatient dermatological care: societal cost-benefit analysis.** *British Medical Journal* 2000, **320**:1252-1256
 38. Rosenfeld BA, Dorman T, Breslow MJ, Pronovost P, Jenckes M, Zhang N, Anderson G, Rubin H: **Intensive care unit telemedicine: alternative paradigm for providing continuous intensivists care.** *Critical Care Medicine* 2000, **28**:3925-3931
 39. Rendina MC: **The effect of telemedicine on neonatal intensive care unit length of stay in very low birthweight infants.** *Proceedings of the 1998 AMIA Annual Fall Symposium* 1998:111-115
 40. Goh KY, Tsang KY, Poon WS: **Does teleradiology improve inter-hospital management of head-injury?** *Canadian Journal of Neurological Sciences* 1997, **24**:235-239
 41. Goh KY, Lam CK, Poon WS: **The impact of teleradiology on the inter-hospital transfer of neurosurgical patients.** *British Journal of Neurosurgery* 1997, **11**:52-56
 42. Lilford RJ, Braunholtz DA, Greenhalgh R, Edwards SJ: **Trials and fast changing technologies: the case for tracker studies.** *British Medical Journal* 2000, **320**:43-46

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright



Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>

editorial@biomedcentral.com